Protostellar collapse: regulation of the angular momentum and onset of an ionic precursor

Pierre Marchand⁽⁰⁾,^{1,2} Kengo Tomida⁽⁰⁾,^{2,3} K. E. I. Tanaka⁽⁰⁾,^{2,4} Benoît Commerçon⁽⁰⁾,⁵ and Gilles Chabrier⁽⁰⁾,^{5,6}

¹American Museum of Natural History, Department of Astrophysics, CPW at 79th, NY, 10024

²Department of Earth and Space Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

³Astronomical Institute, Tohoku University, Sendai, Miyagi 980-8578, Japan

⁴ALMA Project, National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan

⁵CRAL, Ecole normale supérieure de Lyon, UMR CNRS 5574, Université Claude Bernard Lyon 1, France

⁶School of Physics, University of Exeter, Exeter, EX4 4QL, UK

ABSTRACT

Through the magnetic braking and the launching of protostellar outflows, magnetic fields play a major role in the regulation of angular momentum in star formation, which directly impacts the formation and evolution of protoplanetary disks and binary systems. The aim of this paper is to quantify those phenomena in the presence of non-ideal magnetohydrodynamics effects, namely the Ohmic and ambipolar diffusion. We perform three-dimensional simulations of protostellar collapses varying the mass of the prestellar dense core, the thermal support (the α ratio) and the dust grain size-distribution. The mass mostly influences the magnetic braking in the pseudo-disk, while the thermal support impacts the accretion rate and hence the properties of the disk. Removing the grains smaller than 0.1 μ m in the Mathis, Rumpl, Nordsieck (MRN) distribution enhances the ambipolar diffusion coefficient. Similarly to previous studies, we find that this change in the distribution reduces the magnetic braking with an impact on the disk. The outflow is also significantly weakened. In either case, the magnetic braking largely dominates the outflow as a process to remove the angular momentum from the disk. Finally, we report a large ionic precursor to the outflow with velocities of several km s⁻¹, which may be observable.

1. INTRODUCTION

Star formation takes place in molecular clouds when dense regions, the dense cores, undergo gravitational collapse. The regulation of angular momentum (AM) during this process is critical. In particular, the AM is directly linked to the question of planet formation, through the formation of a rotationally-supported protoplanetary disk, and binary star formation by disk fragmentation. Numerous observations (for example Maury et al. 2010) show that the AM is not conserved during the collapse of the dense core and that less than 1% remain in the final star system (Bodenheimer 1995). Early, this phenomenon has been explained by the magnetization of the dense cores (Mestel & Spitzer 1956). Through magnetic braking, magnetic fields extract the AM from the rotating dense core and transports it into the surrounding interstellar medium (ISM).

The first studies including magnetic fields in starformation calculations (both analytical and numerical) were based on ideal magnetohydrodynamics (MHD). In this framework, the magnetic field is perfectly coupled to the gas and undergoes no physical dissipation. The results showed that for realistic magnetizations of dense cores (slightly super-critical), the magnetic braking would remove a significant amount of AM, actually preventing the formation of large disks and strongly hindering fragmentation (Gillis et al. 1974; Matsumoto & Tomisaka 2004; Galli et al. 2006; Price & Bate 2007; Hennebelle & Fromang 2008). This issue is known as the magnetic braking catastrophe. While some amount of turbulence can alleviate the magnetic braking catastrophe (Joos et al. 2012; Santos-Lima et al. 2012; Seifried et al. 2013; Wurster et al. 2019), it became clear that a more accurate description of the magnetic field was necessary and that its decoupling to the gas could not be neglected.

The gas in dense cores is mostly composed of neutral H₂ and He, ~ 70% and ~ 28% by mass, respectively (Draine 2011). The remaining 2 % is a mixture of heavier atoms, molecules and dust grains, either neutral or ionized. The charged species represent only a fraction of 10^{-7} of the particles, but the ideal MHD assumes that the collisional interactions between charged and neutral particles allow the neutrals to couple to the magnetic field whilst not being directly sensitive to the Lorentz's force. More realistically, non-ideal MHD accounts for the decoupling between the magnetic field and neutral species, as well as ions and electrons, through the ambipolar diffusion, the Hall effect and the Ohmic diffusion, respectively. In particular, ambipolar diffusion allows the neutrals to slip through the field lines, reducing the efficiency of magnetic braking. Nowadays most studies use non-ideal MHD, which seems to solve the disk and fragmentation issues of the magnetic braking catastrophe (Machida et al. 2006; Duffin & Pudritz 2008; Mellon & Li 2009; Li et al. 2011; Tomida et al. 2015; Wurster et al. 2016; Masson et al. 2016).

Beside gas, dust grains are also an important component of the ISM. These aggregates of molecules (mostly carbon and silicates) account for 1% of the ISM mass. Because their surface is a catalyst for chemical reactions of gaseous species (e.g. Hocuk & Cazaux 2015), and because they can hold up to several electric charges (Draine & Sutin 1987), they influence the ionisation equilibrium of the gas and the nature of the ionised species. For these reasons, grains play a major role in the coupling of the gas with the magnetic field, thus being a critical factor for non-ideal MHD effects. In the diffuse ISM, the dust grain population is well described by the Mathis, Rumpl, Nordsieck (MRN) size distribution (Mathis et al. 1977), which is commonly used in chemical calculations of non-ideal MHD resistivities (Marchand et al. 2016; Wurster 2016; Koga et al. 2019; Guillet et al. submitted). This distribution can evolve during the protostellar collapse, during which the increasing density promotes the growth of grains by accretion or coagulation (Tielens & Hagen 1982; Rossi et al. 1991; Chokshi et al. 1993; Ormel et al. 2009; Dzyurkevich et al. 2017). These effects can reduce or remove smaller size grains, which is supported by observational evidences (Cardelli et al. 1989; Vrba et al. 1993). Modeling and observing accurate dust size distributions is however a difficult challenge, and assumptions have to be made. Zhao et al. (2016) showed that the removal of smaller grains from the MRN distribution enhances the ambipolar coefficient, which effectively reduces the magnetic braking and helps the formation of large disks. The size distribution of grains is therefore an important parameter of the AM regulation during protostellar collapses.

Another mechanism that regulates the AM are the bipolar outflows, that are widely observed (Cabrit & Bertout 1992; Bontemps et al. 1996; Bacciotti et al. 2002; Tabone et al. 2017) or produced in numerical simulations (Ferreira et al. 2006; Higuchi et al. 2019). While the ejection of low-density fast-rotating gas makes them very efficient for the transport of specific AM, there is however no consensus on whether outflows dominate the magnetic braking to remove the normal AM from the core (see for example Tomisaka 2000; Joos et al. 2012, for two opposite conclusions).

In this study, we analyse the AM regulation in various protostellar collapse simulations with three varying parameters, i.e. the dense core mass, the thermal support and the grain size distribution. The paper is organised as follow. Our methods are described in Section 2, the results are presented in Section 3, including analysis of the disk, outflow and AM transport. We discuss our results in Section 4 and Section 5 is dedicated to conclusion.

2. SIMULATION SETUP

2.1. Theoretical framework

We solve the following MHD equations

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \left[\rho \mathbf{u}\right] = 0, \qquad (1)$$

$$\frac{\partial \rho \mathbf{u}}{\partial t} + \nabla \cdot \left[\rho \mathbf{u} \mathbf{u} + \left(P + \frac{B^2}{2}\right) \mathbb{I} - \mathbf{B} \mathbf{B}\right] = -\rho \nabla \Phi, \quad (2)$$

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \times \left[\mathbf{u} \times \mathbf{B} - \frac{c^2}{4\pi} \eta_{\Omega} \mathbf{J} + \frac{c^2}{4\pi} \eta_{\mathrm{AD}} \frac{(\mathbf{J} \times \mathbf{B}) \times \mathbf{B}}{B^2}\right] = 0(3)$$

$$\nabla \cdot \mathbf{B} = 0. \qquad (4)$$

 ρ is the gas density, **u** is its velocity, *P* is the thermal pressure, **B** is the magnetic field, Φ is the gravitational potential, *c* is the speed of light, η_{Ω} and η_{AD} are the Ohmic and ambipolar diffusion resistivities in s and $\mathbf{J} = \nabla \times \mathbf{B}$ is the electric current. We do not consider the Hall effect in this study. The system is closed by a simple barotropic equation of state (EOS) to mimic the evolution of temperature in collapsing cores (Larson 1969; Masunaga & Inutsuka 2000)

$$T = T_0 \left[1 + \left(\frac{\rho}{\rho_*}\right)^{\gamma - 1} \right], \tag{5}$$

with $T_0 = 10$ K, $\rho_* = 10^{-13}$ g cm⁻³ and $\gamma = 5/3$.

2.2. Initial conditions

Our setup follows the standard Boss & Bodenheimer (1979) initial conditions of a spherical core in solid rotation along the z-axis with an azimuthal density perturbation. We use a rotational energy to gravitational potential energy ratio of $\beta = 0.03$. The density follows a m = 2 azimuthal mode

$$\rho = \rho_0 [1 + \delta_\rho \cos(2\phi)], \tag{6}$$

with ρ_0 the average density, $\delta_{\rho} = 0.1$ the amplitude of the perturbation and ϕ the azimuthal angle. The magnetic field is initially parallel to the rotation axis. Its strength is defined using the mass-to-flux ratio over the critical value

$$\mu = \frac{\frac{M}{\Phi_{\rm B}}}{\left(\frac{M}{\Phi_{\rm B}}\right)_{\rm crit}},\tag{7}$$

where M is the mass of the core, $\Phi_{\rm B} = \pi R_0^2 B_0$ is the magnetic flux, R_0 the initial core radius, B_0 the initial magnetic field strength and

$$\left(\frac{M}{\Phi_{\rm B}}\right)_{\rm crit} = \frac{0.53}{3\pi}\sqrt{\frac{5}{G}} \tag{8}$$

is the critical value, with G the gravitational constant.

The varying parameters are the mass of the core $(M = 2 \text{ or } 5 M_{\odot})$, the thermal to gravitational energy ratio $(\alpha = 0.3 \text{ or } 0.4)$ and the grain size distribution (see Section 2.4). All cases are summarized in Table 1.

In this paper, we refer the "formation time of the first Larson core" $t_{\rm C}$ as the time at which the maximum density reaches 10^{-13} g cm⁻³, and "the first core" as all the gas that exceeds that density. We define the protostellar disk in the same manner as Joos et al. (2012), as the gas verifying the following conditions

-
$$\rho > 3.8 \times 10^{-15} \text{ g cm}^{-3}$$
,

- $u_{\phi} > 2u_r$, with u_{ϕ} and u_r the azimuthal and radial velocities,
- $u_{\phi} > 2u_z$, with u_z the velocity along the z axis,
- $-\frac{\rho u_{\phi}^2}{2} > P$ to ensure that the gas is supported against gravity by its centrifugal force rather than the thermal pressure.

We also define an outflow as the gas verifying the two following conditions

- The velocity is higher than the escape velocity, $||\mathbf{u}|| > u_{\text{lib}} = \sqrt{\frac{2GM_{\text{core}}}{r}}$, where M_{core} is the mass of the first Larson core and r the distance to the center of first core.
- The radial velocity is positive $u_r > 0$.

2.3. Numerical methods

The simulations are performed in 3D using the Adaptive Mesh Refinement (AMR) code RAMSES (Teyssier 2002), incorporating the constrained transport scheme (Londrillo & del Zanna 2004; Fromang et al. 2006; Teyssier et al. 2006) and the ambipolar and Ohmic diffusion (Masson et al. 2012). We use the HLL Riemann solver (Harten et al. 1983) with the minmod slope limiter. The initial grid is uniform and contains 32^3 cells (level 5 of refinement). Cells are then refined to ensure at least 8 points per Jeans length at every location. The stiff equation of state considerably slows down the collapse and prevents the formation of the second Larson core. Considering that the lifetime of the first Larson core is typically a few thousands years (e.g., Vaytet et al. 2013; Bhandare et al. 2018), we stop our simulation 4 to 5 kyr after the formation of the first Larson core. The maximum resolution reached in the simulations is ~ 2 au (level 14 for $M = 2 M_{\odot}$ and level 15 for $M = 5 M_{\odot}$).

2.4. Magnetic resistivities

In our monofluid description, the Ohmic and ambipolar diffusion are controlled by their respective resistivity, which are determined by the chemical environment. We use the table of Marchand et al. (2016), which contains the equilibrium abundances of a reduced chemical network across a wide range of density and temperature. The network includes species relevant to the star formation environment and grains following the MRN size distribution. They account for thermal ionisations, the thermionic emission of grains (Desch & Turner 2015) and the grain evaporation. During the simulation, the resistivities are computed on-the-fly for each cell using the local state variables and tabulated abundances.

For four out of eight simulations, we use another table computed in the same manner with a different grain size distribution. In dense cores, at densities typically higher than the ISM, grains smaller than 0.1 μ m may disappear due to coagulation (Ormel et al. 2009; Guillet et al. submitted). We simulate this coagulation by removing these small grains, while keeping the same dustto-gas ratio as the initial table, $d = 0.0341^{1}$, and the same power-law. Zhao et al. (2016) show that removing grains smaller than 0.1 μ m enhances the ambipolar coefficient the most, hence promoting the formation a large disk due to the reduced magnetic braking. This could be a crucial factor for the AM regulation. The resistivities of both models for the EOS of equation 5 are displayed in Figure 1. Because the temperature increases rapidly, the grain evaporation and the thermal ionizations occur at T > 1000 K, near $\rho \approx 10^{-10}$ g cm⁻³, lowering the resistivities near zero. The gas then becomes perfectly coupled with the magnetic field as in the ideal MHD framework. However, purposely, the increasing thermal support prevents the simulations from reaching higher densities. We qualitatively recover the results of Zhao

¹ In the original table, this dust-to-gas ratio was chosen so that the total surface-area of grains matches the uniform distribution of Kunz & Mouschovias (2009).

Name	Mass (M_{\odot})	α	$ ho_0~({ m g~cm^{-3}})$	R_0 (au)	$\Omega_0 \ (\mathrm{rad} \ \mathrm{s}^{-1})$	$B_0 \ (\mu { m G})$	Grain size distribution
M2a3mrn	2	0.3	1.38×10^{-18}	5900	1.8×10^{-13}	66.7	Standard MRN
M2a3big	2	0.3	1.38×10^{-18}	5900	1.8×10^{-13}	66.7	Truncated MRN
M2a4mrn	2	0.4	5.74×10^{-19}	7900	1.2×10^{-13}	37.5	Standard MRN
M2a4big	2	0.4	5.74×10^{-19}	7900	1.2×10^{-13}	37.5	Truncated MRN
M5a3mrn	5	0.3	2.17×10^{-19}	14830	7.8×10^{-14}	26.7	Standard MRN
M5a3big	5	0.3	2.17×10^{-19}	14830	7.8×10^{-14}	26.7	Truncated MRN
M5a4mrn	5	0.4	9.15×10^{-20}	19770	4.8×10^{-14}	15.0	Standard MRN
M5a4big	5	0.4	9.15×10^{-20}	19770	4.8×10^{-14}	15.0	Truncated MRN

Table 1. Simulation parameters : name of the simulation, mass of the initial sphere, thermal to gravitational energy ratio, average density, radius of the sphere, initial angular velocity, initial magnetic field and grain size distribution.

et al. (2016), with an overall larger ambipolar diffusion and lower Ohmic diffusion coefficients. The main differences is the ambipolar coefficient being smaller in the large-grain cases in the density range $[10^{-14}:10^{-11}]$ $g \text{ cm}^{-3}$, which could affect the dynamics of the first core and the disk. However, we expect the ambipolar diffusion to work against the magnetic braking mostly in the collapsing envelope, at lower densities. Another discrepancy concerns the sign of the Hall effect, that is not affected in our case. However, both their study and ours do not include it in the simulations. The impact of the Hall term on the evolution of the AM has been examined in great detail in Marchand et al. (2018, 2019). Although included in our simulations, the Ohmic diffusion does not play a major role in the dynamics of the disk at these early stages. Its main effect is the dissipation of the magnetic field in the innermost part of the first Larson core. Since the Ohmic resistivity is sensibly the same with both distributions, we will not quantify its effect in this study.

3. RESULTS

Disks and outflows form in every simulation. Their properties are reviewed in Sections 3.1 and 3.2 respectively, while general properties about the transport of AM are discussed in Section 3.3. Section 3.4 is dedicated to the analysis of the ion-neutral drift.

3.1. Disks

Figures 2 and 3 display density maps of the midplane for $M = 2 M_{\odot}$ and $M = 5 M_{\odot}$ respectively, at $t = t_{\rm C} + 1, 2.5$ and 4 kyr (rows 1 to 3), and an edge-on view of the disk (row 4) at $t = t_{\rm C} + 4$ kyr. In our analysis, the mid-plane has a thickness of one cell. The disk selection criteria are described in Section 2.2. The formation of spiral arms, that occurs in all the simulations, is triggered by the initial azimuthal perturbation, and they seem overall more prominent for the 5 solar mass



Figure 1. Non-ideal MHD resistivities as a function of density for the MRN distribution (solid lines), and the truncated distribution (dashed lines). Temperature scales with density according to equation 5, and, for this figure only, we assumed the following magnetic field prescription B = $1.43 \times 10^{-7} \sqrt{\rho/(\mu_{\rm p} m_{\rm H})}$ G (Li et al. 2011), with $\mu_{\rm p} = 2.31$ the mean molecular mass and $m_{\rm H} = 1.67 \times 10^{-24}$ g the mass of one hydrogen atom. The vertical black line represents the first core formation density. $\eta_{\rm H}$ represents the Hall resistivity, which can be negative.

cases and for $\alpha = 0.3$. M2a3mrn, M2a3big, M5a3mrn and M5a4mrn show significant signs of fragmentation. Figures 4 and 5 allow a more quantitative analysis by showing the mass, radius, Toomre's Q and AM of the disks. In table 2, we also summarize the properties of the cores and the disks at $t = t_{\rm C} + 4$ kyr.

We define the disk radius as the cylindrical radius containing 99% the disk's mass. Toomre's Q parameter is often used to quantify the stability of a disk against



Figure 2. Density maps of the four $M = 2 M_{\odot}$ simulations at $t = t_{\rm C} + 1$ kyr (row 1), 2.5 kyr (row 2 and 4) and 4 kyr (row 3). Rows 1 to 3 display slices of the mid-plane, and row 4 shows an edge-on view of the disk at $t - t_{\rm C} + 4$ kyr. In row 4, the gas that does not belong to the disk is not shown.

fragmentation, given by the following formula

$$Q = \frac{c_{\rm s}\Omega_{\rm d}}{\pi G\Sigma},\tag{9}$$

with $c_{\rm s}$ the sound speed in the disk, $\Omega_{\rm d}$ the disk average angular velocity and Σ the surface density of the disk. Q < 1 reflects an instability and a possible fragmentation. The AM is computed over the volume of the disk as

$$\mathbf{L}_{\text{disk}} = \int_{\text{disk}} \rho \mathbf{r} \times \mathbf{u} d\mathbf{V}. \tag{10}$$

In this study, we always show the norm of the AM vector **L**, although its z-component is largely dominant.

Panel a of figures 4 and 5 shows that the disk mass evolution seems to be weakly dependent on the mass of the initial core and the grain size distribution. The low M_{disk} and r_{disk} of M2a4mrn in Table 2 are temporary and due to its fragmentation around this time-mark (see the dark red curve around 4 kyr in the panel a of Figure 5). The higher thermal support of $\alpha = 0.4$ simulations reduces the accretion rate, which yields a lower disk mass. The radii of disks with larger grains are moderately larger for $\alpha = 0.3$, and similar for $\alpha = 0.4$. In the same manner, the Toomre's Q do not largely differ between the two grain models. The slightly larger value for the truncated MRN may be due to a self-regulation of spiral arms formed by gravitational instability, that transfer AM faster when they are more unstable. The difference is more clear when looking at the AM evolution. In the larger grains cases, the disks contain an amount of $\sim 10^{52}$ g cm² s⁻¹ more AM than their MRN cases counter-part, especially after 2.5 kyr, which is an increase of 20% to 50%. The Toomre's Q of all disks are lower than 1 after 2 kyr, which is reflected by the formation of large spiral arms (~ 100 au length). This low value is however not sufficient for the fragmentation of spiral arms, as was noted by Zhao et al. (2018). Takahashi et al. (2016) shows that the spiral arm fragmentation occurs when the local Toomre's Q in the arms becomes lower than 0.6. Therefore, the fragmentation of the disk can not happen as soon as Q < 1, but at later stages as it is the case in our simulations. The



Figure 3. Same as Figure 2 for $M = 5 M_{\odot}$

only run showing no fragmentation by $t = t_{\rm C} + 4$ kyr is M2a4big, which is reflected by the largest Toomre's Q at $t = t_{\rm C} + 4$ kyr. We observe that a larger α provides smaller and less-massive disks with a lower AM. The Toomre's Q is larger and takes more time to reach values below 1. This behavior is mostly a consequence of the lower accretion rate due to the larger free-fall time.

To quantify the influence of the ambipolar diffusion in the disk, we calculate the Elsasser number, which is defined as

$$Am = \frac{4\pi u_A^2}{c^2 \eta_{AD} \Omega},\tag{11}$$

with $u_{\rm A} = B/\sqrt{\rho}$ the Alfven speed, and Ω the angular velocity of the fluid. Am < 1 means that the ambipolar diffusion has a significant impact over the dynamics of the disk. Figure 6 displays the radial profiles of Am in the eight different disks, in which we have performed an azimuthal density-weighted average. Only the innermost 20 to 40 au parts of the disks show Am < 1, down to 10^{-3} . In the outer regions, Am rises up to 10 to 100, indicating a less efficient decoupling between the magnetic field and the gas compared to the central domain. As a consequence of the higher ambipolar resistivity of the reference MRN distribution in the density range $[10^{-14} : 10^{-11}]$ g cm⁻³, the M*a*mrn disks have larger Am < 1 regions than their M*a*big counterpart, up to a factor 2 for M2a3mrn and M2a3big. The M*a*big disks contains however a larger angular momentum (see panel d of figures 4 and 5), meaning that this difference is not significant compared to the influence of the ambipolar diffusion in the outer parts of the disk and in the envelope, although with a higher Elsasser coefficient.

3.2. Outflows

All simulations produce bipolar outflows that start within 1 kyr after the first core formation. Figure 7 shows density maps in the outflow region at $t - t_{\rm C} = 2.5$ kyr. The outflows have similar heart-like structures. It appears that the truncated MRN cases produce outflows with an overall slower expansion and a lower density of a factor ~ 20 ($\approx 5 \times 10^{-17}$ g cm⁻³ vs $\approx 2 \times 10^{-18}$ g cm⁻³) than the standard distribution. They show a clear cavity with dense borders, while cavity only starts to form at 2.5 kyr with the MRN distribution. The weakness of



Figure 4. Time evolution of the mass (panel a), radius (panel b) and Toomre's Q (panel c) and AM (panel d) of the disk, for $\alpha = 0.3$ simulations. Red: $M = 2 M_{\odot}$, blue: $M = 5 M_{\odot}$. Darker colors represent the standard MRN cases while lighter colors represent the truncated MRN.

the truncated MRN outflows are further confirmed when looking at their AM, as shown in Figure 8, that are three to ten times lower than with the standard MRN. The cases with $\alpha = 0.4$ also eject less AM for the normal distribution, but a similar amount than $\alpha = 0.3$ with the truncated MRN. However these quantities do not depend on the dense core's mass. These observations remain qualitatively unaltered if considering the specific AM instead.

In our simulations, the outflows are launched by Blandford & Payne (1982) mechanism. The magnetic field lines are bent by the drag of the collapsing gas, producing the well-known hourglass-shape. If the angle is high enough, the centrifugal acceleration added to the Lorentz force may eject the gas along the field lines. In M^*a^*mrn , the field lines can be bent up to an angle of 75° with the z-axis near the central region, which is a 15° slope for the gas acceleration in the outward radialdirection. However, in M^*a^*big , the higher ambipolar coefficient in the envelope reduces the accretion of the magnetic field due to a weaker coupling with the gas, resulting in a maximum angle of 65°. Figure 9 displays the inclination angle of the magnetic field for both M5a3* simulations, at 2.5 kyr after the formation of the first Larson core, in the plane x = 0 (the pseudo-disk and the contour of the outflow appear clearly). Also due to the decoupling, the magnetic field strength in M*a*big is twice as low as in M*a*mrn in the vicinity of the outflow launching region. Both these effects reduce the magneto-rotational acceleration of the gas.

3.3. Angular momentum transport 3.3.1. The magnetic braking

In this section, we look at the AM regulation in the various simulations. There are two main AM transport mechanisms at play here, that are the advection by the gas (by the gravitational collapse or the outflow), and magnetic braking. To compute the cumulative losses of angular momentum by magnetic braking up to a given time t, we integrate the magnetic torque over the disk or the pseudo-disk



Figure 5. Same as Figure 4 for $\alpha = 0.4$.

Table 2. Summary of the properties of the first cores and disks at $t = t_{\rm C} + 4$ kyr. Mass of the first Larson core, mass of the disk, radius of the disk, Toomre's Q of the disk, mean accretion rate, presence of fragmentation.

Run	$M_{ m C}~(M_{\odot})$	$M_{ m disk}~(M_{\odot})$	$r_{\rm disk}$	Toomre's Q	$\dot{M}_{\rm acc}$	Fragmentation
M2a3mrn	0.30	0.17	123	0.40	7.4×10^{-5}	Yes
M2a3big	0.33	0.19	121	0.40	8.2×10^{-5}	Yes
M2a4mrn	0.19	0.07	52	0.38	4.7×10^{-5}	Yes
M2a4big	0.18	0.13	71	0.52	4.5×10^{-5}	No
M5a3mrn	0.34	0.22	129	0.35	8.5×10^{-5}	Yes
M5a3big	0.34	0.21	143	0.40	8.4×10^{-5}	Yes
M5a4mrn	0.20	0.13	77	0.38	5.1×10^{-5}	Yes
M5a4big	0.20	0.15	82	0.46	5.0×10^{-5}	Yes

$$L_{\text{mag}} = \int_{t_{c}}^{t} \int_{V} r(\mathbf{J} \times \mathbf{B})_{\phi} dV dt.$$
 (12)

where V represents the volume of integration. The criterion to define the pseudo-disk is simply $\rho > 10^{-15}$ g cm⁻³, excluding the disk and the first core as defined in the previous sections.

The profile of the magnetic torque $r (\mathbf{J} \times \mathbf{B})_{\varphi}$ in the mid-plane at $t = t_{\rm C} + 4$ kyr is displayed in Figure 10. The evolutions are similar between the eight cases, with a dispersion across one order of magnitude. The magnetic torque increases as r^2 until $r \approx 100$ au, meaning $(\mathbf{J} \times \mathbf{B})_{\varphi} \propto r$. Then, the torque decreases rapidly as r^{-4} . The location of the peak indicates that the mag-



Figure 6. Radial profiles of the ambipolar Elsasser number (Am) in the disk for the eight simulations at $t = t_{\rm C} + 2.5$ kyr. The quantities have been density-averaged over the azimuthal direction.

netic braking is mostly active in the outer regions of the disk (here $30 \leq r \leq 150$ au). This trend is consistent with the low value of the Elsasser number in the central region, indicating a significant action of the ambipolar diffusion against the magnetic braking. Within the inner 20 au of the disk, the magnetic torque is larger for the standard MRN than for the truncated MRN, by a factor 2 to 10. Conversely, α has a limited impact on the magnetic torque within the innermost 40 au, where the ambipolar diffusion dominates. We observe the opposite phenomenon outside the disk. While the larger α decreases the magnetic torque by one order of magnitude, the grain size-distribution does not alter it. However, we probe here only the mid-plane of the pseudo-disk, in which the ambipolar resistivity is low in both cases. Most of the magnetic braking would take place in outer lavers.

Figure 11 compares the disk and pseudo-disk AM with the amount extracted by magnetic braking, as well as with the AM taken away by the outflow. Panel a) shows that the magnetic braking has removed three to four times (at $t-t_{\rm C} = 5$ kyr) the AM left in the disk, meaning that only 20 to 25 percent of the accreted AM remains in the disk. The magnetic braking is more efficient in the disk for the truncated MRN simulations, which is consistent with a slightly lower ambipolar diffusion coefficient (see Fig. 1). However, panel b) shows that the magnetic braking is the most active in the pseudodisk. With the MRN distribution, more AM is removed by magnetic braking in the pseudo-disk (up to a factor 3 at $t - t_{\rm C} = 5$ kyr). With the truncated distribution, the magnetic braking is similar in both region. The pseudo-disk is therefore of prime importance when

considering the magnetic braking. This is also demonstrated by panel c), which shows the ratio of magnetic braking and AM for the disk and the pseudo-disk combined. Only 10 to 14 percent of the AM remains in this structure for the MRN distribution, and 16 to 25 percent for the truncated MRN cases. The magnetic braking is stronger in the lower-mass cores of M2^{*} simulations than in M5^{*} simulations. Conversely, the α parameter does not change sensitively the amount of AM removed by magnetic braking. Panel d) displays the ratio between the AM taken away by the outflow and the magnetic braking. The magnetic braking largely dominates the outflow by a factor of two to ten with the MRN distribution, and ten to a hundred with the truncated MRN. In this latter case, both the magnetic braking and outflows remove significantly less AM than in the MRN distribution, which further confirms the results of Zhao et al. (2016).

3.3.2. AM in the outflow

Figure 12 shows a slice at the outflow scale of the specific AM, for M5a3mrn. Most of the specific AM of the outflow is located in the bottom wall of the cavity, with values larger than 10^{21} cm² s⁻¹. This area seems connected to the outer layer of the pseudo-disk, that contains a similar specific AM, higher than in the mid-plane at the same cylindrical radius. We have also plotted the projected trajectories of 75 virtual tracer particles, in thin black and white lines. The 50 black particles are randomly generated in the outflow in the final time-step, and the 25 white particles are are generated only in the high specific AM region. In post-processing calculations, we have integrated the particles trajectory backwards using the local flow velocity. This figure shows that the outflowing gas exclusively comes from the envelope and the pseudo-disk. There is almost no direct matter transport from the disk or the mid-plane to the outflow. In the early phases of the outflow, several particles fall near the axis in what we call "the rotor region". This region is a small cylinder of 50-70 au radius and ≤ 10 au height, that is located 50 au above the disk (see the bottom panel). Gas that eventually outflows never goes closer to the disk than the rotor. Particles remain up to 4 kyr in the rotor, rotating one to several times around the axis at $\approx 2 \text{ km s}^{-1}$. There, the radial Lorentz force builds up gradually. The particles eventually move away from the axis and are able to "climb" along a magnetic field line. This occurs when the radial acceleration of the Lorentz force becomes larger than ≈ 0.003 cm s⁻². The specific angular momentum of the particles does not significantly vary when confined in the rotor, and is independent from the time spent in this region. At



Figure 7. Density slices along the plane x=0 for all simulations at $t = t_{\rm C} + 2.5$ kyr. The black lines represent the contour of the outflows and arrows indicate the fluid velocity. Infalling material is marked with black arrows while the outflow is marked with white arrows. Top: M=2 M_{\odot} , bottom: M=5 M_{\odot} . From left to right: M*a3mrn, M*a3big, M*a4mrn, M*a4big.



Figure 8. Time evolution of the AM of the outflow. The curves are the same as in Figure 4 and 5.

later stages, the outflow is fed in both gas and AM by the pseudo-disk, from its sides. In particular, white trajectories in Figure 12 show that the high specific AM region of the outflow is supplied by the outer layers of the pseudo-disk, in which the specific AM is larger than in the mid-plane. A fraction of the collapsing envelope is swept-up by the outflowing gas, which increases the outflow's mass. However, the swept-up gas holds a low AM, thus contributing negatively to the outflow's AM. Additionally, the gas at higher altitudes is launched at an earlier time, coming from more inner regions of the initial cloud, where the specific AM is lower. This explains why the specific AM decreases with height.

3.4. Ion-neutral drift

The ambipolar diffusion is the drift between ions and neutral particles, at the origin of the decoupling between the gas and the magnetic field. The drift velocity is expressed by

$$\mathbf{u}_{\rm AD} = \frac{c^2}{4\pi} \eta_{\rm AD} \frac{\mathbf{J} \times \mathbf{B}}{B^2},\tag{13}$$

and the ion velocity is

$$\mathbf{u}_{\text{ions}} = \mathbf{u} + \mathbf{u}_{\text{AD}},\tag{14}$$

which is also the speed of the magnetic field lines motion because they are attached to the ions in the absence of the Hall effect.

In the mid-plane of the pseudo-disk, due to the pinching of the field lines, B_r rapidly changes with z. For this reason, and because B_z is the main component of the magnetic field, the current is mostly toroidal, so ions are expected to drift in the radial direction (perpendicularly to the current and the magnetic field). The magnetic field lines resist the drag from the gas towards the



Figure 9. Angle between the local magnetic field and the initial magnetic field (vertical) in the y-z plane, at $t = t_{\rm C}+2.5$ kyr. The segments represent the direction of the magnetic field. Top panel: M5a3mrn, bottom panel: M5a3big.

center and tend to "relax" if the decoupling is strong enough. Figure 13 shows the velocity profiles of the gas, the ions and the Keplerian velocity for M5a3mrn (top panel) and M5a3big (bottom panel) in the mid-plane at $t - t_{\rm C} = 4$ kyr. The decoupling is clear in both cases. For M5a3mrn, the ions travel up to 1 km s⁻¹ faster or slower than the neutral gas, and their decoupling is the strongest at the accretion shock, between r = 20to r = 60 au, inside the disk. For r < 20 au, the ions are rotating at a slower speed than the neutral gas. The twisting of the field lines is therefore weaker than what it would be in ideal MHD, reducing the magnetic braking. In M5a3big, between r = 60 and r = 300 au, ions re-



Figure 10. Azimuthal average of the magnetic torque $r (\mathbf{J} \times \mathbf{B})_{\varphi}$ in the mid-plane as a function of the cylindrical radius.

sist the drag from the gas and almost reach a radial and azimuthal velocity near zero between 100 and 200 au. This behavior corresponds to the simulations of Zhao et al. (2018) with a similar grain size distribution. The decoupling in the azimuthal direction is not as strong overall, the velocity difference reaching half the values of the radial direction.

We now look at the ion-neutral decoupling at larger scales, in the vicinity of the outflow. Figure 14 displays the relative velocities of the ions and the neutrals for M5a3mrn and M5a3big at $t - t_{\rm C} = 2.5$ kyr. The color represents the drift velocity $||\mathbf{u}_{AD}|| = ||\mathbf{u} - \mathbf{u}_{ions}||$ in the z and ϕ directions, while the white and black arrows indicate the direction of the neutrals and the ions respectively. Inside the pseudo-disk, the drift velocity is negligible, indicating a strong coupling, similarly to the velocity profiles in Figure 13 beyond few 100 au. In the infalling envelope however, the ions weakly decouple as they approach the mid-plane, until they enter the pseudo-disk. For $|z| \lesssim 500$ au, their trajectory is more bent toward the center than the neutrals. In the outflow of M5a3big, the drift velocity reaches several 10 km s⁻¹, similar to the value of the Alfven velocity, due to the extremely high ambipolar coefficient (two to four orders of magnitude higher than in the envelope or the disk). However, the most striking features are the large zones of strong decoupling in the vicinity of the outflow for the truncated MRN case. Due to the high ambipolar diffusion, caused by the low density, and the strong electric current and magnetic field close to the central axis, the ions drift from the neutrals at velocities of several km s^{-1} . We even observe an outflow of ions outside and bigger than the actual previously-



Figure 11. Panel a: ratio between the cumulative magnetic braking in the disk and the AM of the disk. Panel b: ratio between the cumulative magnetic braking in the pseudo-disk (not including the disk) and the AM of the disk. Panel c: Ratio between the cumulative magnetic braking in the disk + pseudo-disk and the AM of the disk + pseudo-disk. Panel d: Ratio between the AM transported by the outflow and the cumulative magnetic braking in the disk.

described outflow. Ions are escaping against the flow of infalling neutrals from the envelope at velocities of several km s⁻¹. The velocity difference between both kind of species exceeds 5 km s⁻¹ in these regions. The ions also strongly drift azimuthally at velocities of typically 2 km s⁻¹. From now, to distinguish both structures, we call this decoupling region the "ion-outflow", and the previously defined outflow as the "neutral-outflow". To our knowledge, such behavior has never been reported in actual star forming clouds observations nor in simulations. In the surrounding infalling envelope, ions also drift from the neutrals, though at lower speed, and seem to converge towards the ion-outflow in the same way that the neutral gas converges toward the neutral-outflow, as

shown in Figure 12. This ion-outflow is actually a nonideal magnetosonic wave driven by the neutral outflow, that is supersonic (Mach number $\mathcal{M} = u/c_{\rm s} > 5$) but sub-Alfvenic ($\mathcal{M}_{\rm a} = u/c_{\rm A} < 0.5$). This wave propagates in the surrounding medium and accelerates ions ahead of the neutrals because of their much higher Alfvén speed. This phenomenon is countered by the ion-neutral drag, that is much higher in the standard MRN case, because a larger grain surface area leads to more collisions, which prevents the formation of a large precursor ion-outflow.

4. DISCUSSION

The question of the regulation of AM is deeply related to the formation of binary stars, particularly, in



Figure 12. Specific AM of the gas in the outflow region in the plane x = 0, for M5a3mrn. The thicker black line represent the boundaries of the outflow at $t - t_{\rm C} = 5$ kyr. Thin lines are the trajectories, projected onto the yz-plane, of virtual tracer particles located in the outflow in the last output, and whose path is integrated backward in time. The 50 black particles are randomly distributed in the outflow, and the 25 white particles are distributed only in the region of high specific AM. The brown triangles indicate the location of the particles in the last output. For clarity, the abscissa of white trajectories represent the cylindrical radius and not the actual y-coordinate. The red countour represents the disk. The bottom panel is a zoom of the top panel.

our non-turbulent case, by disk fragmentation. Machida et al. (2008) performed an extensive study on this subject by varying the magnetic field strength and the rotational velocity. They were able to define parameter ranges suitable for the formation of wide binary systems (separation of 3-300 au), close binary systems (0.007-0.3 au) and single stars. The value of our parameters falls within the wide binary region (see their figure 12), which is consistent with the high instability of the disks in our simulations (spiral arms > 100 au, low Toomre's Q,



Figure 13. Velocity profiles in the mid-plane for M5a3mrn (top panel) and M5a3big (bottom panel) at $t - t_c = 4$ kyr. Red curves represent the radial velocities, blue curves are the azimuthal velocities and the black curve indicate the Keplerian velocity. Solid lines represent the gas while dashed lines represent the ions.

fragmentation). The major differences are their initial condition, a Bonnor-Ebert sphere (Ebert 1955; Bonnor 1956), and the absence of ambipolar diffusion, with only the Ohmic diffusion. Nonetheless, in our study, a uniform sphere yields a higher accretion rate and the ambipolar diffusion weakens the magnetic braking, which promote the fragmentation and the formation of wide binaries. With a longer evolution, we could expect the formation of stable wide companions by disk fragmentation. In their 3D simulations of protostellar collapse with ambipolar diffusion, Hennebelle et al. (2020) found that the properties of the disk are weakly dependent on the initial conditions, except the magnetization but including the initial rotation velocity. This is a result that we concur. The disks are however heavily influenced by the accretion scheme on sink particles, which we do



Figure 14. Relative velocities of the ions and the neutrals for M5a3mrn (left panels) and M5a3big (right panels) at $t - t_{\rm C} = 2.5$ kyr. The color represents the drift velocity $||\mathbf{u}_{\rm AD}|| = ||\mathbf{u} - \mathbf{u}_{\rm ions}||$ along the z-direction (top panels) and ϕ -direction (bottom panels). White arrows : neutral gas velocity direction, black arrows : ions velocity direction, red contour : neutral gas outflow.

not have in this study. The presence of a sink particle reduces the mass and the extent of the disk, hence its ability to fragment.

The relative importance between the magnetic braking and the outflow as an AM removal mechanism is still debated. The early model of Tomisaka (2000) shows a dominance of the outflow over the magnetic braking in what would typically be the first Larson core phase. This work used the ideal MHD framework. Our study shows that ambipolar diffusion weakens the outflow even more than the magnetic braking. Therefore, the outflow may be able to dominate the magnetic braking in the absence of ambipolar diffusion. On the other hand, using 3D ideal MHD simulations, Joos et al. (2012) finds that the magnetic braking transports more AM than the outflow. Depending on the orientation between the magnetic field and the rotation axis, the two effects are quantitatively closer than in our study, which confirms our hypothesis on the results of Tomisaka (2000). In our simulations, we find that at least 75% of the AM accreted in the central region is removed by magnetic braking, even with high ambipolar coefficients. We can therefore estimate that the outflows (including later protostellar jets) cannot account for more than 25% of the AM transport in the presence of ambipolar and Ohmic diffusion. At later stages, in an already formed disk without accretion, Pudritz & Ray (2019) argue that the outflow should be the dominant process of angular momentum transport, because the Magneto-Rotational Instability (MRI) is greatly reduced by non-ideal MHD effects (Blaes & Balbus 1994; Bai & Stone 2013). Additionally, outflows driven by MRI can be significantly enhanced by the presence of the Hall effect (Lesur et al. 2014).

In Section 3.4 we report for the first time an ionoutflow precursor to the neutral gas outflow. With a typical ionisation of 10^{-7} , the ion density in this region would be only of the order of $\approx 10 \text{ cm}^{-3}$. However, with a velocity difference of several km s^{-1} between ions and neutral, observations should be able to detect the precursor, using the CO molecule, a common tracer in protostellar outflows, and HCO⁺, one of the most abundant ion and also a common tracer in young star systems. Yen et al. (2018) performed an observation of the ion-neutral drift in the Class 0 protostar B335 with the Atacama Large Millimeter Array (ALMA) telescope. They did not detect any strong decoupling down to the 100 au scale, finding an upper limit of 0.3 km s⁻¹ on the drift velocity in the collapsing gas. The absence of a significant decoupling would indicate a weaker ambipolar diffusion than in our simulations, which is confirmed by the small size of the disk, less than 10 au. Since a Class 0 protostar is more advanced than the systems we present in this study, another possibility is that the ion-outflow is a short-lived phenomenon that has already disappeared. In any case, more observations of the ion-neutral drift are needed to bring constraints on the chemistry at stake, and most importantly the grain size-distribution, and as a proof of the importance of non-ideal MHD.

There are limitations to the monofluid approximation that could affect the ion-outflow. The inertia of charged particles is neglected, meaning that the velocity of ions always immediately reaches the theoretical velocity calculated from the drift velocity. Additionally, ions could be subject to a chemical drag: charge transfer reaction between the outflowing ions and the infalling neutrals would effectively reduce the average velocity of the ion fluid. It is difficult to quantify these effects, but it is possible that they reduce the drift velocity. More accurate results necessitate a bifluid framework and chemical calculations on-the-fly. We discuss this subject more extensively in appendix A.

5. CONCLUSIONS

We have performed 3D simulations of protostellar collapses to study the regulation of AM through the analysis of the disk, the outflow and the magnetic braking. Our parameters are the mass and the thermal support of the initial sphere of gas, and the grain-size distribution, either the MRN distribution or a truncated MRN distribution from which small grains have been removed. Our main results are the following.

- The mass of the dense core has little impact on the properties of the disk and the outflow during the first Larson core phase.
- A higher thermal over gravitational energy ratio α reduces the accretion rate which yields smaller and lower-mass discs, and reduces the AM transported by the outflow in the MRN case, but not in the truncated MRN case.
- The influence of the mass of the dense core and α on the magnetic braking in the disk is extremely limited, supporting the idea that the first core and the disk "forget" some of the initial conditions.
- Removing the small grains produces slightly larger disks containing more angular momentum due to the higher ambipolar diffusion resistivity, but the difference is not significant. The outflow is however considerably weaker, transporting 10 times less AM compared to the MRN distribution.
- Most of the gas and AM of the outflow comes from the outer layers of the pseudo-disk.
- By the end of the first core phase, the magnetic braking is responsible for up to 80 percent of AM loss in the disk, and 75 percent (for the truncated MRN) to 90 percent (for the MRN distribution) in the disk + pseudo-disk.
- While not negligible, the transport of AM by the outflow is largely below the magnetic braking, by a factor 5 (for the MRN distribution) to 100 (for the truncated MRN).
- In the truncated MRN simulations, the high ambipolar diffusion leads to an ion-outflow that is a precursor of the neutral-outflow. Ions travel at several km s⁻¹ against the infalling envelope, at larger scale than the neutral outflow. Observational evidences of such structures would be informative regarding the chemistry at stake in protostellar collapses.

The conclusions show that non-ideal MHD plays a significant role in the regulation of AM during the protostellar collapse, and that the results heavily depend on the chemical model, especially the grain-size distribution.

ACKNOWLEDGMENTS

The authors thank Kazunari Iwasaki and Mordecai-Mark Mac-Low for their thoughts and advices that greatly helped our analysis. We acknowledge financial support from an International Research Fellowship of the Japan Society for the Promotion of Science (JSPS), number P17802, from the "Programme National de Physique Stellaire" (PNPS) of CNRS/INSU, CEA and CNES, France, and from the American Museum of Natural History. Computations were performed on the computer Occigen (CINES) under the allocation DARI A0020407247 made by GENCI, and on the Center For Computational Astrophysics of the National Astronomical Observatory of Japan, under a XC-B allocation. K. Tomida was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers 16H05998, 16K13786, 17KK0091, 18H05440. K.Tanaka acknowledges support from NAOJ ALMA Scientific Research Grant 2017-05A, JSPS KAKENHI Grant JP19H05080, JP19K14760.

APPENDIX

A. ION-NEUTRAL COLLISION RATES

In this section we analyse the influence of the drift velocity onto the ambipolar resistivity, because we deem important to emphasize some points regarding the calculation of the non-ideal MHD resistivities in the one-fluid approximation. The detail of the formulae can be found in several studies (Nakano & Umebayashi 1986; Marchand et al. 2016). Collision rates between ions and neutral are needed in the calculation, and most recent studies (Marchand et al. 2016; Wurster 2016; Zhao et al. 2016) take these numbers from Pinto & Galli (2008). The fitting formulae of their Table 2 are commonly used, and depend on the ion velocity

$$v_{\rm rms} = \sqrt{v_{\rm d}^2 + \frac{8kT_{ss'}}{\pi\mu_{ss'}}},$$
 (A1)

where v_d is the drift velocity between species s and s', k the Boltzman constant, $T_{ss'}$ the weighted average temperature, that we consider to be the fluid temperature in the one-fluid approximation, and $\mu_{ss'}$ the reduced mass of both species. In multi-fluid calculations, it is possible to derive v_d from the various species velocities. Since the quantity is not directly accessible in the one-fluid framework, it is often assumed to be zero. However, in cases of strong ambipolar drift, the drift velocity may very well exceed the thermal velocity, the second term. In this case, the approximations on collision rates, and resistivities, would prove inaccurate. Moreover, as detailed in their table 2, the fitting formulae of momentum transfer coefficients between ions and an H₂ are not valid if $v_{\rm rms}$ exceeds 5 to 10 km s⁻¹. We therefore have to be extremely careful in our analysis. In this paper, the drift velocity we compute in our simulations may exceed 10 km s⁻¹, but always in the neutral-outflow. We therefore consider our results about the ion-neutral drift inaccurate om this region.

We use the drift velocity v_d as a parameter and look at its influence on the ambipolar resistivity. We assume the same drift velocity for all molecular ions. Small grains (< 1 μ m) being well coupled to the gas (see for instance Laibe & Price 2014; Lebreuilly et al. 2019), we do not expect them to significantly drift from H₂. Electrons have a high thermal velocity, their $v_{\rm rms}$ is thus not strongly affected by a drift velocity of a few km s⁻¹. Here, we have assumed T = 10 K, and $B = 1.43 \times 10^{-7} \sqrt{\rho/(m_{\rm p}\mu_{\rm p})}$ (Li et al. 2011). Ambipolar resistivities as a function of the drift velocity are displayed in Figure 15, for various densities, and for both grain size distributions.



Figure 15. Ambipolar resistivity as a function of v_d for various densities. Red color: $\rho = 10^{-18}$ g cm⁻³, blue color: $\rho = 10^{-16}$ g cm⁻³, purple color: $\rho = 10^{-14}$ g cm⁻³, green color: $\rho = 10^{-12}$ g cm⁻³. The solid lines represent the MRN case and the dashed lines represent the truncated MRN case.

Below $v_d = 1 \text{ km s}^{-1}$, the ambipolar coefficient does not seriously diverges from its value at $v_d = 0$. However, the difference is significant at higher velocities for the truncated MRN, while the MRN case shows a more reduced variation. At 10 km s⁻¹, the resistivities are larger by a factor 2 to 5 for the truncated MRN, and extends beyond one order of magnitude at 1000 km s⁻¹ (although it is unlikely that such high speed would be reached in the context of star formation). This difference comes from the fact that grains are the dominant charged species in the MRN case, while it is the ions in the truncated MRN case, because of the lower number of grains in this density range. As a consequence, with a truncated MRN, the resistivities are more affected by variation of the ions drift velocities. In the outflow, the density is of the order of $\rho = 10^{-18} \text{ g cm}^{-3}$ and the magnetic field $B \approx 0.01 \text{ G}$, which is higher than the value given by our prescription, but the results are qualitatively the same.

Collapsing the ions and neutrals momentum and energy equations removes one degree of freedom in the system, as the ambipolar resistivity represents itself the drift between both fluids. This issue represents a limit of the monofluid description of non-ideal MHD.

REFERENCES

- Bacciotti, F., Ray, T. P., Mundt, R., Eislöffel, J., & Solf, J. 2002, ApJ, 576, 222, doi: 10.1086/341725
- Bai, X.-N., & Stone, J. M. 2013, ApJ, 769, 76, doi: 10.1088/0004-637X/769/1/76
- Bhandare, A., Kuiper, R., Henning, T., et al. 2018, ArXiv e-prints. https://arxiv.org/abs/1807.06597
- Blaes, O. M., & Balbus, S. A. 1994, ApJ, 421, 163, doi: 10.1086/173634
- Blandford, R. D., & Payne, D. G. 1982, MNRAS, 199, 883, doi: 10.1093/mnras/199.4.883
- Bodenheimer, P. 1995, ARA&A, 33, 199, doi: 10.1146/annurev.aa.33.090195.001215
- Bonnor, W. B. 1956, MNRAS, 116, 351
- Bontemps, S., Andre, P., Terebey, S., & Cabrit, S. 1996, A&A, 311, 858

- Boss, A. P., & Bodenheimer, P. 1979, ApJ, 234, 289, doi: 10.1086/157497
- Cabrit, S., & Bertout, C. 1992, A&A, 261, 274
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245, doi: 10.1086/167900
- Chokshi, A., Tielens, A. G. G. M., & Hollenbach, D. 1993, ApJ, 407, 806, doi: 10.1086/172562
- Desch, S. J., & Turner, N. J. 2015, ApJ, 811, 156, doi: 10.1088/0004-637X/811/2/156
- Draine, B. T. 2011, Physics of the Interstellar and Intergalactic Medium
- Draine, B. T., & Sutin, B. 1987, ApJ, 320, 803, doi: 10.1086/165596
- Duffin, D. F., & Pudritz, R. E. 2008, MNRAS, 391, 1659, doi: 10.1111/j.1365-2966.2008.14026.x

- Dzyurkevich, N., Commerçon, B., Lesaffre, P., & Semenov,
 D. 2017, A&A, 603, A105,
 doi: 10.1051/0004-6361/201628995
- Ebert, R. 1955, Zeitschrift für Astrophysik, 37, 217
- Ferreira, J., Dougados, C., & Cabrit, S. 2006, A&A, 453, 785, doi: 10.1051/0004-6361:20054231
- Fromang, S., Hennebelle, P., & Teyssier, R. 2006, A&A, 457, 371, doi: 10.1051/0004-6361:20065371
- Galli, D., Lizano, S., Shu, F. H., & Allen, A. 2006, ApJ, 647, 374, doi: 10.1086/505257
- Gillis, J., Mestel, L., & Paris, R. B. 1974, Ap&SS, 27, 167, doi: 10.1007/BF00641596
- Guillet, V., Hennebelle, P., Marcowith, A., et al. submitted
- Harten, A., Lax, P., & Van Leer, B. 1983, SIAM Review, 25, 35, doi: 10.1137/1025002
- Hennebelle, P., Commerçon, B., Lee, Y.-N., & Charnoz, S. 2020, A&A, 635, A67, doi: 10.1051/0004-6361/201936714

Hennebelle, P., & Fromang, S. 2008, A&A, 477, 9, doi: 10.1051/0004-6361:20078309

- Higuchi, K., Machida, M. N., & Susa, H. 2019, MNRAS, 486, 3741, doi: 10.1093/mnras/stz1079
- Hocuk, S., & Cazaux, S. 2015, A&A, 576, A49, doi: 10.1051/0004-6361/201424503
- Joos, M., Hennebelle, P., & Ciardi, A. 2012, A&A, 543, A128, doi: 10.1051/0004-6361/201118730
- Koga, S., Tsukamoto, Y., Okuzumi, S., & Machida, M. N. 2019, MNRAS, 484, 2119, doi: 10.1093/mnras/sty3524
- Kunz, M. W., & Mouschovias, T. C. 2009, ApJ, 693, 1895, doi: 10.1088/0004-637X/693/2/1895
- Laibe, G., & Price, D. J. 2014, MNRAS, 440, 2136, doi: 10.1093/mnras/stu355
- Larson, R. B. 1969, MNRAS, 145, 271, doi: 10.1093/mnras/145.3.271
- Lebreuilly, U., Commerçon, B., & Laibe, G. 2019, A&A, 626, A96, doi: 10.1051/0004-6361/201834147
- Lesur, G., Kunz, M. W., & Fromang, S. 2014, A&A, 566, A56, doi: 10.1051/0004-6361/201423660
- Li, Z.-Y., Krasnopolsky, R., & Shang, H. 2011, ApJ, 738, 180, doi: 10.1088/0004-637X/738/2/180

Londrillo, P., & del Zanna, L. 2004, Journal of Computational Physics, 195, 17, doi: 10.1016/j.jcp.2003.09.016

- Machida, M. N., Inutsuka, S., & Matsumoto, T. 2006, ApJL, 647, L151, doi: 10.1086/507179
- Machida, M. N., Tomisaka, K., Matsumoto, T., & Inutsuka, S.-i. 2008, ApJ, 677, 327, doi: 10.1086/529133
- Marchand, P., Commerçon, B., & Chabrier, G. 2018, A&A, 619, A37, doi: 10.1051/0004-6361/201832907
- Marchand, P., Masson, J., Chabrier, G., et al. 2016, A&A, 592, A18, doi: 10.1051/0004-6361/201526780

- Marchand, P., Tomida, K., Commerçon, B., & Chabrier, G. 2019, A&A, 631, A66, doi: 10.1051/0004-6361/201936215
- Masson, J., Chabrier, G., Hennebelle, P., Vaytet, N., & Commercon, B. 2016, A&A, 587, A32, doi: 10.1051/0004-6361/201526371
- Masson, J., Teyssier, R., Mulet-Marquis, C., Hennebelle, P., & Chabrier, G. 2012, ApJS, 201, 24, doi: 10.1088/0067-0049/201/2/24
- Masunaga, H., & Inutsuka, S. 2000, ApJ, 531, 350, doi: 10.1086/308439
- Mathis, J. S., Rumpl, W., & Nordsieck, K. H. 1977, ApJ, 217, 425, doi: 10.1086/155591
- Matsumoto, T., & Tomisaka, K. 2004, ApJ, 616, 266, doi: 10.1086/424897
- Maury, A. J., André, P., Hennebelle, P., et al. 2010, A&A, 512, A40, doi: 10.1051/0004-6361/200913492
- Mellon, R. R., & Li, Z. 2009, ApJ, 698, 922, doi: 10.1088/0004-637X/698/1/922
- Mestel, L., & Spitzer, Jr., L. 1956, MNRAS, 116, 503
- Nakano, T., & Umebayashi, T. 1986, MNRAS, 218, 663, doi: 10.1093/mnras/218.4.663
- Ormel, C. W., Paszun, D., Dominik, C., & Tielens,
 A. G. G. M. T. 2009, A&A, 502, 845,
 doi: 10.1051/0004-6361/200811158
- Pinto, C., & Galli, D. 2008, A&A, 484, 17, doi: 10.1051/0004-6361:20078819
- Price, D. J., & Bate, M. R. 2007, Ap&SS, 311, 75, doi: 10.1007/s10509-007-9549-x
- Pudritz, R. E., & Ray, T. P. 2019, Frontiers in Astronomy and Space Sciences, 6, 54, doi: 10.3389/fspas.2019.00054
- Rossi, S. C. F., Benevides-Soares, P., & Barbuy, B. B. 1991, A&A, 251, 587
- Santos-Lima, R., de Gouveia Dal Pino, E. M., & Lazarian, A. 2012, ApJ, 747, 21, doi: 10.1088/0004-637X/747/1/21
- Seifried, D., Banerjee, R., Pudritz, R. E., & Klessen, R. S. 2013, MNRAS, 432, 3320, doi: 10.1093/mnras/stt682
- Tabone, B., Cabrit, S., Bianchi, E., et al. 2017, A&A, 607, L6, doi: 10.1051/0004-6361/201731691
- Takahashi, S. Z., Tsukamoto, Y., & Inutsuka, S. 2016, MNRAS, 458, 3597, doi: 10.1093/mnras/stw557
- Teyssier, R. 2002, A&A, 385, 337, doi: 10.1051/0004-6361:20011817
- Teyssier, R., Fromang, S., & Dormy, E. 2006, Journal of Computational Physics, 218, 44, doi: 10.1016/j.jcp.2006.01.042
- Tielens, A. G. G. M., & Hagen, W. 1982, A&A, 114, 245
- Tomida, K., Okuzumi, S., & Machida, M. N. 2015, ApJ, 801, 117, doi: 10.1088/0004-637X/801/2/117
- Tomisaka, K. 2000, ApJL, 528, L41, doi: 10.1086/312417

- Vaytet, N., Chabrier, G., Audit, E., et al. 2013, A&A, 557, A90, doi: 10.1051/0004-6361/201321423
- Vrba, F. J., Coyne, G. V., & Tapia, S. 1993, AJ, 105, 1010, doi: 10.1086/116489
- Wurster, J. 2016, PASA, 33, e041, doi: 10.1017/pasa.2016.34
- Wurster, J., Bate, M. R., & Price, D. J. 2019, MNRAS, 489, 1719, doi: 10.1093/mnras/stz2215
- Wurster, J., Price, D. J., & Bate, M. R. 2016, MNRAS, 457, 1037, doi: 10.1093/mnras/stw013
- Yen, H.-W., Zhao, B., Koch, P. M., et al. 2018, A&A, 615, A58, doi: 10.1051/0004-6361/201732195
- Zhao, B., Caselli, P., Li, Z.-Y., & Krasnopolsky, R. 2018, MNRAS, 473, 4868, doi: 10.1093/mnras/stx2617
- Zhao, B., Caselli, P., Li, Z.-Y., et al. 2016, MNRAS, 460, 2050, doi: 10.1093/mnras/stw1124